

SUSY 2013, 26-31 August 2013, Trieste

Top quark production in the ATLAS detector of the LHC

Javier Montejo Berlingen, in behalf of the ATLAS collaboration

Instituto de física de altas energías, IFAE



- Top has a large production cross section. Including associated production because of the large phase space.
- Complex final state, top pair production is background for a number of SUSY searches

0 leptons + 2 b-jets + E _{miss}	[S _{bottom} /stop]	arxiv:1308.2631
0 leptons + ≥ 7 -10 jets + E _{miss}	[Incl. strong production]	arxiv:1308.1841
2 leptons + (b)jets + E _{miss}	[Medium stop]	ATLAS-CONF-2013-065
1-2 leptons + 3-6 jets + E _{miss}	[Incl. strong production, mUED]	ATLAS-CONF-2013-062
0-1 leptons + ≥ 3 b-jets + E _{miss}	[3rd gen. Squarks]	ATLAS-CONF-2013-061
1 lepton + 4(1 b-)jets + E _{miss}	[Medium / heavy stop]	ATLAS-CONF-2013-037
0 lepton + 6 (2 b-)jets + E _{miss}	[Heavy stop]	ATLAS-CONF-2013-024
1-2 taus + jets + E _{miss}	[GMSB]	ATLAS-CONF-2013-026

- The LHC has produced a large dataset of tops to analyze
 - This allows for differential measurements, observation of difficult decay channels, associated production, etc
- Precision measurements allow for a better understanding and reduced systematics
- I will show results from six ATLAS results, showing inclusive and differential measurements

- Top quark is the heaviest SM particle, decays ~100% of the times to Wb
- Analyses are defined by decay of W boson

Single lepton (e/μ) + jets	34.3%
Dilepton ($ee/e\mu/\mu\mu$)	6.4%
All hadronic	45.7%
Tau + jets	9.8%
Tau + lepton	3.7%

- 2 b-quarks per event, usage of b-tagging to increase $t\bar{t}$ contribution

Top Pair Decay Channels

$c\bar{s}$	electron+jets	muon+jets	tau+jets	all-hadronic		
$u\bar{d}$						
τ^-						
μ^-	$e\mu$	$\mu\mu$	$\mu\tau$	muon+jets		
e^-	ee	$e\mu$	$e\tau$	electron+jets		
W decay	e^+	μ^+	τ^+	$u\bar{d}$		$c\bar{s}$

- **Total inclusive cross section**

- Measurement of the top quark pair production cross section in the single-lepton channel with ATLAS in proton-proton collisions at 8 TeV using kinematic fits with b-tagging: [Link](#)
- Measurement of the $t\bar{t}$ production cross section in the tau+jets channel using the ATLAS detector: [Link](#)

- **Differential cross section**



- Measurement of $t\bar{t}$ production with a veto on additional central jet activity in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector: [Link](#)
- Measurement of the jet multiplicity in top-anti-top final states produced in 7 TeV proton-proton collisions with the ATLAS detector: [Link](#)
- Measurements of top quark pair relative differential cross-sections with ATLAS in pp collisions at $\sqrt{s} = 7$ TeV: [Link](#)

- **Associated production**

- A study of heavy flavor quarks produced in association with top quark pairs at $\sqrt{s} = 7$ TeV using the ATLAS detector: [Link](#)

Top pair cross section in lepton+jets

- Dataset: 5.8 fb^{-1} @8TeV
- 1 lepton (e/μ , $p_T > 40 \text{ GeV}$), 3 or more jets ($p_T > 25 \text{ GeV}$), 1 or more b-tags (70% efficiency), large E_T^{miss} and $m_T(W)$
- Likelihood discriminant based on transformed aplanarity^[*] and lepton pseudorapidity
- Fit the discriminant for $t\bar{t}$ and W+jets normalization
→ extract $t\bar{t}$ cross section



• Result: $\sigma_{t\bar{t}} = 241 \pm 2(\text{stat}) \pm 31(\text{syst}) \pm 9(\text{lumi}) \text{ pb}$

$$\text{NNLO+NNLL: } \sigma_{t\bar{t}}^{\text{NNLO}} = 253_{-15}^{+13} \text{ pb}$$



[*] $A' = \exp(-3/4 \lambda_3)$

λ_3 being the smallest eigenvalue of the normalized momentum tensor

ATLAS-CONF-2012-149

Top pair cross section in tau+jets

- Dataset: 1.67 fb^{-1} @7TeV
- 5 or more jets ($p_T > 20 \text{ GeV}$), 2 or more b-tags (60% efficiency), high E_T^{miss} significance
- Top candidate is built from 1 tagged + 2 untagged jets with highest p_T sum.
Remaining non b-tagged jet is τ candidate
- Likelihood fit to number of tracks distribution with three templates: τ /electron, gluon-jet (from multi-jet), quark-jet (from $t\bar{t}$ and W +jets)
- Result: $\sigma_{t\bar{t}} = 194 \pm 18(\text{stat}) \pm 46(\text{syst}) \text{ pb}$
NNLO+NNLL: $\sigma_{t\bar{t}}^{\text{NNLO}} = 177_{-11}^{+10} \text{ pb}$
- Why tau+jets?
 - Probe flavor-dependent effects, test lepton universality
 - Background for BSM scenarios such as charged Higgs





- Dataset: 2.05 fb^{-1} @7TeV, dilepton channel
- Very precise measurement. Is able to constrain modeling uncertainties and reduce their impact on other top quark measurements
- Select dilepton events with 2 b-tags, very high purity

- Define gap fraction:
$$f(Q_0) = \frac{n(Q_0)}{N}$$



N is the number of selected events, $n(Q_0)$ is the subset that do **not** contain an additional central jet with $p_T > Q_0$

- Compare NLO generators (MC@NLO, POWHEG), multileg LO generators (ALPGEN, SHERPA) and initial state radiation systematics (ISR) with AcerMC, and constrain the allowed range for the variations.
- All generators predict too much forward activity. MC@NLO predicts too little activity in central region






- Dataset: 4.7 fb^{-1} @7TeV, lepton+jets channel
- Motivation: constrain models of additional QCD radiation in $t\bar{t}$ events, test perturbative QCD
- Differential measurement in N_{jets} for jet p_T thresholds at 25, 40, 60, 80 GeV
- Background-subtracted events, unfolded into particle-jet multiplicity distribution
- MC@NLO+HERWIG model disfavored, predicts lower jet multiplicity and softer jets
- ALPGEN+PYTHIA nominal and α_s -up predict too high jet multiplicities

[*] Stable particles except muons and neutrinos, clustered with anti-kt algorithm, $R=0.4$



Top pair differential cross section

- Dataset: 2.05 fb^{-1} @7TeV, lepton+jets channel
- Relative differential cross-sections as a function of the invariant mass $M_{t\bar{t}}$, the transverse momentum $p_{T,t\bar{t}}$, and the rapidity $y_{t\bar{t}}$ of the top quark pair system
- Precision SM test. $M_{t\bar{t}}$ sensitive to new physics in s-channel
- 1 lepton (e/μ , $p_T > 25 \text{ GeV}$), 4 or more jets ($p_T > 25 \text{ GeV}$), 1 or more b-tags (70% efficiency), large E_T^{miss} and $m_T(W)$
- Kinematic likelihood fit of the event, cut on the likelihood improves purity of the sample
- Background-subtracted events, unfolded into parton-level kinematic distribution
- Measurement consistent with SM expectation





- Dataset: 4.7 fb^{-1} @7TeV, dilepton channel
- Fiducial cross section measurement and ratio, 2 charged leptons ($p_T > 25 \text{ GeV}$, $|\eta| < 2.5$), 3 or more jets ($p_T > 25 \text{ GeV}$, $|\eta| < 2.5$)
- $t\bar{t}$ +HF measured from events with at least 3 b-tags
 $t\bar{t}$ +j measured from events with 3 jets, 2 b-tags
- Maximum likelihood fit to separate light/heavy flavor, 2D templates: vertex mass and jet p_T
- Fit performed in 3 mutually exclusive bins of b-jet purity. Boundaries defined at operating points of 60, 70, 75% efficiency

$$R_{\text{HF}} = \frac{\sigma_{\text{fid}}(t\bar{t} + \text{HF})}{\sigma_{\text{fid}}(t\bar{t} + \text{j})} = 7.1 \pm 1.3(\text{stat})_{-2.0}^{+5.3}(\text{syst})\%$$

Compared to ALPGEN 3.4% and POWHEG 5.2%

- Why $t\bar{t}$ +HF?
 - Background for $t\bar{t}H(H \rightarrow bb)$, $H \rightarrow tb$, composite Higgs, 4 tops



- Results agree well with SM theory predictions
- ATLAS is testing the SM at high precision with inclusive and differential cross section measurements
 - In some cases the measurements are reaching enough precision to constrain the range of systematic variations and disfavor generator models
 - Smaller uncertainties → better sensitivity for searches :)
- Better understanding of QCD in $t\bar{t}$ final states through differential measurements paves the way for more precision in the future
- Please see: <https://twiki.cern.ch/twiki/bin/view/AtlasPublic/TopPublicResults> for the full list of ATLAS top quark publications, only a subset has been shown here