The μ problem and the NMSSM with strings

Saúl Ramos-Sánchez
UNAM

SUSY 2013

Aug. 27, 2013

In collaboration with

R. Kappl, M. Ratz, K. Schmidt-Hoberg, P. Vaudrevange; and O. Lebedev

Top-Down approach: From strings to the real world

Top-Down approach: From strings to the real world

Top-Down approach: From strings to the real world



Top-Down approach: From strings to the real world



$E_8 \times E_8$ heterotic orbifolds

Heterotic String en 10D



input: Orbifold	output: effective QFT in 4D
Geometry	gauge symmetry $\mathcal{G}_{4D}$
Embedding	matter spectrum
($\mathbb{Z}_N$, Lattice(s), Twist, Shifts, Wilson lines, discrete torsion)	further symmetries $\rightarrow K,\ W,\ f_a$

$E_8 \times E_8$ heterotic orbifolds

Heterotic String en 10D



In this talk...

- get close to (N)MSSM
- solve some hierarchy issues

From minilandscape and further searches

☺ there are "fertile patches"

Our search:

- ~ 300 "MSSM"
- no charged exotics
- grand unification
- heavy top
- $m_{3/2} \sim$TeV
- seesaw
- R parity
- flavor symmetries
- QCD axion
- $\mu H_u H_d \; with \; \mu \ll M_{Pl}$

From minilandscape and further searches

☺ there are "fertile patches"

Our search:

- ~ 300 "MSSM"
- no charged exotics
- grand unification
- heavy top
- $m_{3/2} \sim$TeV
- seesaw
- R parity
- flavor symmetries
- QCD axion
- $\mu H_u H_d \; with \; \mu \ll M_{Pl}$

$\left.\begin{array}{l}\text{no charged exotics}\\ \text{grand unification}\\ \text{heavy top}\end{array}\right\}$ required

From minilandscape and further searches

☺ there are "fertile patches"

Our search:

- ~ 300 "MSSM"
- no charged exotics
- grand unification
- heavy top
- $m_{3/2} \sim$TeV
- seesaw
- R parity
- flavor symmetries
- QCD axion
- $\mu H_u H_d \; with \; \mu \ll M_{Pl}$

$\left.\begin{array}{l}\text{no charged exotics}\\ \text{grand unification}\\ \text{heavy top}\end{array}\right\}$ required

$\left.\begin{array}{l}m_{3/2} \sim \text{TeV}\\ \text{seesaw}\\ R \text{ parity}\\ \text{flavor symmetries}\\ \text{QCD axion}\\ \mu H_u H_d \; with \; \mu \ll M_{Pl}\end{array}\right\}$ gift !! ☺

From minilandscape and further searches

☺ there are "fertile patches"

Our search:

- ~ 300 "MSSM"
- no charged exotics
- grand unification
- heavy top
- $m_{3/2} \sim$TeV
- seesaw
- R parity
- flavor symmetries
- QCD axion
- $\mu H_u H_d$ *with* $\mu \ll M_{Pl}$?

required

gift !! ☺

Hierarchies

from

R symmetries

$\mathrm{U}(1)_R$ symmetries

- Perturbative superpotential

$$\mathscr{W} \;=\; \sum c_{n_1 \cdots n_M} \phi_1^{n_1} \cdots \phi_M^{n_M}$$

- vacuum with $\mathcal{N} = 1$

$$-F_i^\dagger \;=\; \frac{\partial \mathscr{W}}{\partial \phi_i} \;=\; 0 \quad \text{in } \phi_j = \langle \phi_j \rangle \;\forall\; i, j$$

- $\mathrm{U}(1)_R$ symmetry

$$\mathscr{W} \;\to\; \mathrm{e}^{2\mathrm{i}\,\alpha}\, \mathscr{W} \qquad\qquad \phi_j \;\to\; \phi_j' \;=\; \mathrm{e}^{\mathrm{i}\, r_j\, \alpha}\, \phi_j$$

$$\mathscr{W}(\phi_i) \;\to\; \mathscr{W}(\phi_i') \;=\; \mathscr{W}(\phi_i) + \sum_j \underbrace{\frac{\partial \mathscr{W}}{\partial \phi_j}}_{=0} \Delta\phi_j$$

$$\Rightarrow \mathscr{W} \;=\; 0$$

$\mathrm{U}(1)_R$ symmetries

$\mathrm{U}(1)_R \quad \& \quad F = 0 \quad \Longrightarrow \quad$ vacuum with $\langle \mathscr{W} \rangle = 0$

Consequences:

- $D\mathscr{W} = 0$ in sugra $\quad \rightarrow \quad \langle V \rangle = 0$: Minkowski vacuum ☺
- does not matter if $\mathrm{U}(1)_R$ is exact or accidental ☺
- complements Nelson+Seiberg's theorem:

$\mathscr{W}$ with $\mathrm{U}(1)_R \quad \Longrightarrow$

- SUSY vacuum with $\langle \mathscr{W} \rangle = 0$
- SUSY or non-SUSY vacuum

Nelson, Seiberg (1994)

accidental $\mathrm{U}(1)_R$

If $\mathrm{U}(1)_R$ is accidental, i.e. explicitly broken at order N:

1. $\langle \mathscr{W} \rangle \sim \langle \phi \rangle^{\geq N}$
2. $\mathscr{W}_{eff} = \langle \mathscr{W} \rangle + \mathscr{W}_{np}$
3. $m_{3/2} \sim \langle \mathscr{W} \rangle$
4. Goldstone mode η gets mass $m_\eta \sim m_{3/2}/\langle \phi \rangle^2$

If e.g. $\langle \phi \rangle \sim \mathcal{O}(0.1)$:

- $N \sim \mathcal{O}(10) \quad \Rightarrow \quad$ hierarchically suppressed $m_{3/2}$ ☺
- η a bit heavier than the gravitino ☺

In string models

In heterotic orbifolds...

No global stringy $\mathrm{U}(1)$s allowed, but...



accidental symmetry $\mathrm{U}(1)_R$ broken at order $N \in [10, 20]$ ☺

Magnitude of $\langle \phi_i \rangle$ determined by

$$D_{\mathrm{FI}} = \xi + \sum_i q^i_{\mathrm{anom}} |\langle \phi_i \rangle|^2 \stackrel{!}{=} 0\,, \;\; \xi \sim \mathcal{O}(0.1) \text{ in Planck units}$$

$$\rightarrow \quad \langle \phi_i \rangle \sim \mathcal{O}(0.1)$$

Assume moduli stabilization here!

In heterotic orbifolds...

- $H_u H_d \supset \mathbf{1}$ (gauge and stringy symmetries)

Antoniadis, Gava, Narain, Taylor; Brignole, Ibáñez, Muñoz; Cvetič, Louis, Ovrut

$$\rightarrow \; K \supset -\log[(T+\bar{T})(U+\bar{U}) - (H_u+\bar{H}_d)(H_d+\bar{H}_u)]$$

$$\rightarrow \; \mu \sim \langle \mathscr{W} \rangle$$

Brümmer, Kappl, Ratz, Schmidt-Hoberg

In SUSY vacua

$$\langle \mathscr{W} \rangle \sim \mathcal{O}(10^{-20}) - O(10^{-10}) \quad \rightarrow \quad \mu \; suppressed! \; ☺$$

Assume moduli stabilization here!

Stringy NMSSM

NMSSM

MSSM + singlet S, such that

$$W = W_{MSSM} + \lambda S H_u H_d + \frac{1}{3}\kappa S^3$$

$$V(s) = -2\lambda A_\lambda v_u v_d s + m_S^2 s^2 + (\kappa s^2 - \lambda v_u v_d)^2 + (\lambda v_d s)^2 + \frac{2}{3}\kappa A_\kappa s^3$$

E.g. in the PQ limit (slightly) broken ($\kappa \ll 1$)



$$\rightarrow \quad \mu = \lambda\langle s\rangle \gtrsim 100 \text{ GeV}$$

Heterotic NMSSMs

In heterotic orbifolds

- μ term *highly* suppressed
- one (or <u>very</u> few) massless S charged under other gauge $\mathrm{U}(1)$s
- renormalizable coupling $S\,H_u\,H_d$
- S^3 possible only at (large) non-renormalizable orders

$$\Rightarrow \quad \lambda = \mathrm{cte.} + \langle s_{a_1} ... s_{a_n} \rangle, \qquad \kappa = \langle s_{b_1} ... s_{b_n} \rangle$$

NMSSMs heteróticos: generalidades

Magnitude of $\langle s_i \rangle$ given by

$$D_{\rm FI} = \xi + \sum_i q^i_{\rm anom} |\langle s_i \rangle|^2 \stackrel{!}{=} 0\,,\ \xi \sim \mathcal{O}(0.1)$$

$$\rightarrow \quad \langle s_i \rangle \sim \mathcal{O}(0.1)$$

$$\rightarrow \quad \boxed{\kappa \ll 1, \qquad \lambda \lesssim \mathcal{O}(1)} \qquad ☺$$

Two possible NMSSM scenarios:

- $\lambda \sim 1,\ \kappa \ll 1$ PQ limit
- $\lambda,\ \kappa \ll 1$ "effective MSSM"

NMSSMs heteróticos: generalidades

Magnitude of $\langle s_i \rangle$ given by

$$D_{\rm FI} = \xi + \sum_i q^i_{\rm anom} |\langle s_i \rangle|^2 \stackrel{!}{=} 0\,,\ \xi \sim \mathcal{O}(0.1)$$

$$\rightarrow \quad \langle s_i \rangle \sim \mathcal{O}(0.1)$$

$$\rightarrow \quad \boxed{\kappa \ll 1, \qquad \lambda \lesssim \mathcal{O}(1)}$$

☺

Two possible NMSSM scenarios:

- $\boxed{\lambda \sim 1,\ \kappa \ll 1 \qquad \text{PQ limit}}$
- $\lambda,\ \kappa \ll 1$ "effective MSSM"

NMSSMs heteróticos: generalidades

Magnitude of $\langle s_i \rangle$ given by

$$D = \xi + \sum_i q^i_{\text{anom}} |\langle s_i \rangle|^2 \stackrel{!}{=} 0\,,\ \ \xi \sim \mathcal{O}(0.1)$$

$$\rightarrow \quad \langle s_i \rangle \sim \mathcal{O}(0.1)$$

$$\rightarrow \quad \boxed{\kappa \ll 1, \qquad \lambda \lesssim \mathcal{O}(1)} \qquad ☺$$

In a sample heterotic model with a SUSY vacuum

$$W_{NMSSM} = S H_u H_d + \tfrac{1}{3} \langle s_i \rangle^6 S^3$$

In this and most heterotic NMSSM models

$\lambda \sim 1,\ \ \kappa \sim 10^{-6}\ \rightarrow$ límite con $m_{a_{PQ}} \sim 10^2$ MeV – GeV ☺

To take home...

In orbifolds with other appealing properties

- mecanism to yield hierarchies based on $\mathrm{U}(1)_R$ ✓
- consequence: $\mu \ll M_{Pl}$ ✓

- NMSSM less common than MSSMs
- S^3 coupling suppressed

  $\rightarrow$ PQ limit or "effective MSSM" ✓
- solutions to μ & fine-tuning problems of MSSM ✓
- R & PQ symmetries come from string theory (*not ad hoc*) ✓